"Utilizing Time Domain Reflectometry to Investigate Dielectric Behavior in Binary Polar Liquid Mixtures"

Synopsis

Submitted To

Swami Ramanand Teerth Marathwada University, Nanded

For The Registration of
Ph. D. Degree
In Physics, Faculty of Science and Technology
Swami Ramanand Teerth Marathwada University, Nanded

Submitted By
Mr. Venkatesh Hemchandra Vispute
M.Sc.
School of Physical Science S.R.T.M.U. Nanded.

Under The Supervision of
Dr. B.G. Nemmaniwar
Assistant Professor & HOD
Department of Physics
Digambarrao Bindu ACS College Bhokar Dist.Nanded

1. Introduction

The intricate dance of molecules within the realm of chemistry has long captivated the inquisitive minds of scientists leading to the unraveling of profound mysteries that shape our understanding of the material world. Among the myriad facets of molecular interactions, dielectric properties stand as a fundamental yet intricate domain offering insights into the electrostatic behavior and structural intricacies of matter. In this pursuit of knowledge we embark on a scientific odyssey that endeavors to explore the dielectric behavior of amines chemical entities renowned for their diverse applications in both aqueous solutions. Dielectric measurements grounded in the science of dielectric spectroscopy have emerged as a potent tool for probing the electrical properties of materials across a wide frequency spectrum. These measurements unlock a treasure trove of information about the polarization, dipole dynamics and charge distribution within substances. From elucidating the fundamental principles of salvation and molecular interactions to facilitating advancements in fields as diverse as chemistry, biology and materials science, the significance of dielectric studies transcends disciplinary boundaries [1-7].

The Enigmatic Role of amines at the heart of our investigation lies a class of compounds known as amines. These molecules characterized by the presence of amino ($-NH_2$) groups are imbued with unique properties that render them indispensable in numerous industrial, biological and chemical processes. Amines both aromatic and aliphatic have carved a niche in applications ranging from pharmaceuticals and polymers to agrochemicals and corrosion inhibition [8].

The behavior of molecules within solvents is a realm of profound interest and the choice of solvent can dramatically influence their dielectric behavior. Aqueous solutions with their propensity for hydrogen bonding and a high dielectric constant exhibit distinct characteristics where a diverse landscape of dielectric behaviors awaits exploration. In this endeavor, we chart a comprehensive research program aimed at unraveling the dielectric mysteries of amines in aqueous solvents. Our journey encompasses the meticulous measurement of dielectric parameters including the static dielectric constant, dielectric constant at high frequency, relaxation time and the elusive Kirkwood correlation factor. Through a symphony of experimental rigor, data analysis and theoretical modeling we aspire to illuminate the structural and dynamic aspects of amines in diverse solvent environments.

As we embark on this scientific voyage we anticipate that our research will not only contribute to the understanding of dielectric phenomena but also hold implications for fields as diverse as chemistry, biology, materials science and environmental science. The journey of discovery awaits,

promising to uncover the hidden melodies of molecular interactions and the intricate dance of amines in the realm of dielectric measurements.

2. Review of literature and development in the subject

Certainly, let's delve into a brief review of the literature and developments in the subject of dielectric measurements and the behavior of amines in different solvents. Dielectric Relaxation Spectroscopy (DRS) is a powerful technique used to investigate the electrical properties of materials as a function of frequency. Dielectric relaxation spectroscopy is occupied very large range of frequency band i.e. from 10^4Hz to 10^{12}Hz [9]. Dielectric relaxation spectroscopy is sensitive to relaxation process in an extremely wide range of characteristic time i.e. 10^5-10^{-12} seconds. By using this spectroscopy we can understand the liquid dynamics charge distribution, orientation of molecules and solute-solvent interaction within binary and ternary mixtures. It has been widely employed to study the dielectric behavior of various substances including polar and non-polar solvents. Amine-Solvent Interactions the interaction of amines with solvents is a topic of significant interest due to its relevance in various fields such as chemistry, biology and materials science. Amines being polar molecules can form hydrogen bonds with solvent molecules leading to changes in their dielectric properties [10-13]. Static dielectric constant (ε_0), the static dielectric constant of a substance reflects its ability to store electrical energy under a static electric field. It is a fundamental parameter that characterizes the polarity and the charge distribution within a material [14]. Dielectric relaxation time (τ), dielectric relaxation time measures the time taken by dipoles in a material to respond to changes in an external electric field. It is a crucial parameter for understanding the dynamics of polar molecules in solvents [15]. Dielectric behavior in aqueous solutions dielectric studies in aqueous solutions have revealed distinct behaviors. Aqueous solutions often exhibit higher dielectric constants due to the strong hydrogen bonding interactions between water molecules. Kirkwood correlation factor (g), Kirkwood correlation factor is a measure of the local molecular order within a liquid. It provides insights into the degree of structuring and clustering of molecules in a solution [9]. Recent developments, recent advancements in dielectric spectroscopy have seen the application of advanced techniques such as time-domain dielectric relaxation spectroscopy (TDRS) and broadband dielectric spectroscopy. These techniques allow for more comprehensive characterization of dielectric properties over a wide frequency range. Applications understanding the dielectric behavior of amines in different solvents has practical applications in various fields. For example, it can inform the design of solvents for chemical reactions aid in the development of pharmaceuticals and contribute to the study of molecular interactions in biological systems [16]. Modeling and theoretical approaches theoretical models such as the Debye Model and the Havriliak-Negami equation mentioned in your proposed work, have been used to interpret dielectric data and gain insights into molecular dynamics and interactions [17]. Environmental and industrial implications, dielectric studies can also have implications for environmental monitoring and industrial processes. Understanding how solvents and chemicals behave electrically can aid in pollution control and process optimization [18-22].

In summary, the literature in this field spans a wide range of topics from fundamental dielectric properties to their practical applications. Our proposed research aims to contribute to this body of knowledge by studying the dielectric behavior of amines in different solvents which has relevance in multiple scientific and industrial domains.

3. Objective Research / Proposed Hypothesis

In the realm of scientific exploration our grand endeavor takes flight, soaring into the vast expanse of knowledge. Behold the proposed hypothesis that shall guide our noble quest! In pursuit of enlightenment, we shall embark on a journey of meticulous investigation. Our mission, none less than the probing of amines secrets submerged in aqueous realms by employing the venerable Time Domain Dielectric Relaxation Spectroscopy technique. A majestic spectrum of frequencies spanning from the mellifluous 10MHz to the resounding crescendo of 50GHz, shall be our compass through this uncharted terrain. With steely determination our quest unfurls its banners revealing its noble objectives.

First, we shall unravel the enigmatic tapestry of **static dielectric constant**, **complex permittivity and relaxation time** that shrouds these amines within the aqueous embrace. Time Domain Reflectometry, our trusted ally shall wield its powers to unveil this mysterious veil. In our grasp we shall seize the elusive dielectric constant, the spectral dance of the **dielectric loss** and the rhythmic heartbeat of relaxation time all resounding at the exalted frequencies of the high realm.

But our aspirations reach even higher for we yearn to delve into the very soul of these molecules. Their interactions shall be scrutinized with the precision of a virtuoso's performance. Behold the metrics of our scrutiny, the **Kirkwood correlation factor, the Bruggman Factor and Excess Parameters** all bow before the might of our Picoseconds Dielectric Relaxation Spectroscopy.

In sum, our journey shall be one of a we and revelation where the arcane world of amines in aqueous media yields its secrets and the symphony of high-frequency dielectric measurements resonates through the annals of scientific discovery. With flourish and fervor we embark on this noble odyssey!

4. Methodology to be adopted

In our quest for knowledge, a meticulous methodology shall serve as our compass with precision akin to an artisan's craft we shall employ the following steps to unravel the complex dielectric spectra through the venerable Time Domain Reflectometry (TDR) technique.

Experimental Setup. Our journey commences with the Tektronix model no. DSA8300 Digital Serial Analyzer, a paragon of precision and its accompanying sampling module. This tandem shall wield the powers of Time Domain Reflectometry. Behold Figure 1, which unveils the experimental setup testament to our scientific endeavor [23]. **Pulse Generation.** In the heart of our experiment lies a rhythmic heart beata repetitive, fast-rising 250mV voltage pulse, a symphony resonating at 200 KHz. Its pulse has a swift 12ps incident and a reflective rise time of 15ps. It courses through a coaxial line system with an impedance of 50 ohms, a conductor of our inquiries [24]. **Sampling Oscilloscope.** As our pulse travels the vigilant eye of a sampling oscilloscope stands sentinel. It captures the metamorphosis of the pulse as it reflects from the end of its journey. The reflected pulse snaked and veiled by the sample bear witness to our experiment. A 5ns time window is their canvas and they are painted in the strokes of 2000 points as depicted in Figure 2. **Data Manipulation.** The pulses dance in pairs one from the pristine reflection (R1(t)) and the other entwined with the essence of the sample (Rx(t)). They are summoned forth both in their original form [q(t)=R1(t) + RX(t)] and in their contrast [p(t) = R1(t) - RX(t)]. These forms reveal the essence of our investigation [25].

Transformation into the Frequency Domain. With the data in hand we embark on a voyage from the time domain into the realm of frequencies. The Fourier transformation a magician's incantation shall conjure the spectral secrets hidden within. Summation and Samulon methods shall be our loyal companions in this endeavor. Our goal, to reach frequencies up to 50GHz and therein to uncover the frequency-dependent complex reflection coefficient spectra, $\rho^*(\omega)$, as per [9].

$$(\rho)^*=cp\ (\omega)/j\ \omega d\ q(\omega) \qquad (1)$$

where c is velocity of light in mm/ps, ω is angular frequency, d is effective pin length. The complex permittivity spectra $\varepsilon^*(\omega)$ was obtained from complex reflection spectra $\rho^*(\omega)$ by using bilinear calibration method given by

The Quest for Complex Permittivity: The elusive complex permittivity spectra $\varepsilon^*(\omega)$, await our discovery. From the complex reflection spectra $\rho^*(\omega)$ we shall extract this enigma using the bilinear calibration method a method encapsulated by the equation [26].

$$\varepsilon^*(\omega)=\varepsilon_{\infty+}(\varepsilon_0-\varepsilon_\infty)/[(1_+(j\omega\tau)^{1-\alpha}]^{\beta} \qquad (2)$$

Here, ε_0 stands as the static dielectric constant, ε represents the permittivity at high frequency, τ is the relaxation time, while α and β embody the symphony of symmetric and asymmetric distribution of relaxation times, respectively. Behold the Havriliak-Negami equation portal into the mysteries of three relaxation models each with its own unique character.

1. If $\alpha = 0$ and $\beta = 1$, we embrace the simplicity of a single Debye relaxation.
2. When $0 \leq \alpha \leq 1$, we traverse the labyrinth of the Cole-Cole model symmetric distribution of relaxation times.
3. And when $\alpha = 0$, while β ranges from 0 to 1 we encounter the enigmatic Cole-Davidson (CD)an asymmetric distribution of relaxation times.

With these meticulous steps and the symphony of our instrumentation, we embark on a journey to reveal the hidden facets of complex dielectric spectra an odyssey that shall enlighten the corridors of scientific understanding.



(a)

(b)

Fig1. (a)Block diagram of Time Domain Reflectometry. (b) Photograph of TDR



Fig.2 Reflected pulse without sample *R1(t)* & sample *Rx(t)*.

5.Importance of Study/Society application

Aniline and its halogenated derivatives in combination with alkoxyalcohols have diverse applications across several industries due to their unique properties. Some of these applications include.

Pharmaceuticals. Aniline derivatives are fundamental building blocks in pharmaceutical synthesis contributing to the production of various medicines and drugs. Halogenated anilines may possess specific pharmacological properties useful in drug development. Alkoxyalcohols might serve as solvents or intermediates in pharmaceutical formulations. **Dyes and Pigments.** Aniline and its derivatives are crucial in the manufacturing of dyes and pigments used in textiles, plastics and other materials. The introduction of halogenated groups can modify the color spectrum and properties of these dyes. alkoxyalcohol could be utilized as solvents or stabilizers in dyes production. **Polymers and Plastics.** Aniline derivatives contribute to the production of various polymers and plastics. The incorporation of halogenated groups

can alter the mechanical, thermal and chemical properties of these materials. alkoxyalcohol might function as plasticizers, aiding in improving flexibility or processability of polymers.

The applications of aniline and its halogenated derivatives in combination with alkoxyalcohol highlight their versatility and importance across a wide range of industries, owing to their distinctive properties and reactivity enabling their use in various processes and formulations.

6. International status/National status:

1. S. Yaghara, University of Tokyo, Japan.
2. N. Shiyanshiki, University of Tokyo, Japan.
3. S. Sudo, dept. of Physics, Tokyo city University, Tamazustumi, Setagaya, Tokyo.
4. A. Catldo, university di Lecce, via Monteroni, 73100 Lecce, Italy.
5. S.C. Mehrotra, Dr. BAMU, Aurangabad(MS).
6. A. C. Kumbharkhane, SRTM University, Nanded (MS).
7. R.J. Sengwa, Jodhpur, JNV University, Jodhpur(RS).

7. Proposed work plan

Proposed work plan for a three-year research project on dielectric measurements of haloanilines and substituted alcohols seems well-structured. Here's a breakdown of the plan.

First Year: Literature Survey and Solvent Selection

- **Objective:** To build a strong foundation for my research and select appropriate solvents for our experiments.
- **Activities:**
  - Conduct an extensive literature survey to understand the existing knowledge in the field.
  - Identify key research gaps and areas where our study can contribute.
  - Familiarize our self with dielectric measurement techniques and equipment.
  - Begin the process of solvent selection, considering factors like solubility, stability and relevance to my research.

Second Year: Dielectric Measurements

- **Objective:** To conduct dielectric measurements on haloanilines with substituted alcohols and extract essential dielectric parameters.
- **Activities:**
  - Acquire and set up the necessary equipment for dielectric measurements.
  - Prepare samples of amines in various solvents as per my solvent selection criteria.
  - Perform dielectric measurements at different frequencies.
  - Extract dielectric parameters including static dielectric constant, dielectric constant at high frequency, relaxation time, Bruggeman factor, Kirkwood correlation factor, excess properties.

Third Year: Dielectric and Structural Behavior Study

- **Objective:** To delve deeper into the dielectric and structural behavior of haloanilines with substituted alcohols by analyzing dielectric relaxation parameters.
- **Activities:**
  - Analyze the data collected from dielectric measurements.
  - Study how dielectric parameters vary with changes in concentration, temperature or other relevant factors.
  - Investigate the structural aspects and molecular interactions within the solutions.
  - Determine any correlations or trends between dielectric behavior and the molecular structure of the haloanilines with substituted alcohols.
  - Summarize your findings and draw conclusions regarding the dielectric and structural behavior of haloanilines in substituted alcohols.

Throughout the three years, it's crucial to maintain meticulous records of my experiments, data analysis and observations. Additionally consider incorporating regular progress assessments, presentations and discussions with our research advisor or team to ensure that my research stays on track and aligns with my objectives.

References

[1] T. Koltai, M. Macaud, A. Guevara, E. Schulz, M. Lemaire, R. Bacaud, M. Vrinat, Applied Catalysis A, General 231, 2002, 253–261.

[2] K.H. Choi, Y. Sano, Y. Korai, I. Mochida, Applied Catalysis B Environmental, 53, 2004, 275–283.

[3] L. Vradman, M.V. Landau, M. Herskowitz, Catalysis Today, 48, 1999, 41–48.

[4] A. Ishihara, T. Itoh, T. Hino, M. Nomura, P.Y. Qi, T. Kabe, Journal of Catalysis, 140, 1993,184–189.

[5] X. Zhang, D.O. Hayward, M.P. Mingos, Catalysis Letters, 88, 2003, 1–2.

[6] J. Chen, M. Te, H. Yang, Z. Ring, Petroleum science and technology, 21, 5 & 6, 2003, 911-935.

[7] N.K. Mehrotra, S.B.L. Misra, S.K.S. Somevanshi, Pramana, 11, 1978, 187–190.

[8] B. G. Nemmaniwar, N. V. Kalyankar,P. L. Kadam, Orbital Electron Journal Chemistry, 2013, 5, 1-6.

[9] S. Sudo, N. Oshiki, N. Shinyashiki, Journal Physics Chemistry, A, 111,16, 2007, 2993-2998.

[10] F. Kremer, A. Schonhals. Broadband Dielectric Spectroscopy, Springer, 2003.

[11] B.G. Nemmaniwar,A.I.P.ConferenceProceeding, 2269, 2020,030114-1–030114-4.

[12] B.G. Nemmaniwar,Materials Today Proceedings, 23, 2020, 324–328.

[13] B. G. Nemmaniwar, Journal of Chemical, Biological and Physical Sciences, Section C, 12, 3, 2022,202-211.

[14] S.C. Mehrotra, A.C. Kumbharkhane& A.S. Chaudhari. Binary Polar Liquids,Elsevier, 2017.

[15] N.K. Karthick, G. Arivazhagan, A.C. Kumbharkhane, Journal of Molecular Structure, 1108, 2016, 203-208.

[16] R.H. Cole, J.G. Berberian, S. Mashimo, G. Chryssikos, A. Burns, E. Tomb, Journal of Applied Physics, 66, 1989, 793–802.

[17] D. Bertolini, M. Cassettari, G. Salvetti, E. Tomabari, S. Versoni, The Review of Science, Instrctrtion, 61, 1990,450–456.

[18] A.C. Kumbharkhane, S.M. Puranik, S.C. Mehrotra, FaranceTransation, 87,1991. 1569–1573.

[19] S. Havriliak, S. Negami, Journal of Polymer Science Part C, 14, 1966,99–117.

[20] P. Debye, Polar Molecules, the Chemical Catalogue Company, New York, 1929.

[21] K.S. Cole, R.H. Cole, The Journal of Chemical Physics, 9, 1941,341–345.

[22] D.W. Davidson, R.H. Cole, The Journal of Chemical Physics, 18, 1950, 1484–1487.

[23] A. Karunakaran, T. Ganesh, M Maria Sylvester, Fluid Phase Equilibria, 382, 2014, 1-17.

[24] Y. S. Joshi, K. S. Kanse, D. N. Rander, A. C. Kumbharkhane, Indian Journal Pure & Applied Physics, 54, 2016, 621-628.

[25] R. J. Sengwa, S. Sankhla, N. Shinyashiki, Journal Solution Chemistry, 37, 2008, 137-153.

[26] A. C. Kumbharkhane, Y. S. Joshi, S. C. Mehrotra, S. Yagihara, S. Sudo, Physica B 421, 2013, 1-7.

Mr. Veankatesh H. Vispute
Research Student
School of Physical Science S.R.TM.U.
Nanded.

Dr. B.G. Nemmaniwar
Research Supervision
Assistant Professor & HOD
Department of Physics
Digambarrao Bindu ACS College
Bhokar. Dist. Nanded.

RESEARCH SUPERVISOR
Digambarrao Bindu ACS College
Bhokar Dist.Nanded